



GRS 1943-052

Lund, Niels; Castro-Tirado, A. J.; Brandt, Søren

Published in:
International Astronomical Union Circulars (IAUC)

Publication date:
1991

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Lund, N., Castro-Tirado, A. J., & Brandt, S. (1991). GRS 1943-052. *International Astronomical Union Circulars (IAUC)*, (5370). <http://www.cbat.eps.harvard.edu/iauc/05300/05370.html#Item2>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Central Bureau for Astronomical Telegrams
 INTERNATIONAL ASTRONOMICAL UNION
 Postal Address: Central Bureau for Astronomical Telegrams
 Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.
 Telephone 617-495-7244/7440/7444 (for emergency use only)
 TWX 710-320-6842 ASTROGRAM CAM EASYLINK 62794505
 MARS DEN or GREEN@CFA.BITNET MARS DEN or GREEN@CFAPS2.SPAN

RS OPHIUCHI

Y. Andrillat and L. Houziaux, Observatoire de Haute Provence, communicate: "Spectra of the recurrent nova RS Oph have been obtained during Aug. 17-19 in the ranges 759-800, 837-878, and 982-1020 nm at 0.1-nm resolution. The broad and violet-shifted emission wings mentioned by Ijima ([IAUC 5302](#)) are no longer observed. All lines are narrow (0.3-0.5 nm FWHM). Monochromatic magnitudes in the continuum are 9.45 (787 nm), 10.47 (850 nm), 10.36 (880 nm), and 9.54 (1 micron), somewhat brighter than at minimum light. The following fluxes ($\times 10^{-12}$ erg cm $^{-2}$ s $^{-1}$) of Fe II emission lines were observed: 771.1 nm, 0.42; 786.8 nm, 1.98; 999.7 nm, 0.36; these fluxes have been corrected for $E(B-V) = 0.8$. The brightest line is O I at 844.6 nm (2.63×10^{-12} erg cm $^{-2}$ s $^{-1}$). The Paschen decrement is very mild, since from P7 to P17 the flux varies from 1.8 to 0.47×10^{-12} erg cm $^{-2}$ s $^{-1}$. The Ca II triplet is quite conspicuous, but no forbidden lines are detected."

GRS 1943-052

N. Lund, A. J. Castro-Tirado, and S. Brandt, on behalf of the Granat WATCH team (Danish Space Research Institute, Lyngby; and Space Research Institute, Moscow), report: "A short-duration x-ray transient, GRS 1943-052 (R.A. = 19h43m, Decl. = -5.2, equinox 1950.0, error radius 1 deg), was recorded by the WATCH all-sky x-ray monitor on Oct. 16.48 UT. The x-ray flux rose in 20 s, reaching a peak flux of about 10 times that of the Crab Nebula in the 6- to 15-keV energy band, and thereafter decayed below the detection limit in 50 s. Follow-up observations are encouraged."

SUPERNOVA 1991bd IN UGC 2936

J. Mueller reports her independent discovery of this object (cf. [IAUC 5367](#)), which was at mag about 16.5 on a blue plate exposed by J. D. Mendenhall and herself with the 1.2-m Oschin Telescope on Oct. 15 UT during the course of the second Palomar Sky Survey. A spectrogram obtained by M. Mateo and I. Reid on Oct. 16 with the 5-m Hale Telescope (+ double spectrograph) suggests that this is a type-II supernova.

1991 October 18

(5370)

Daniel W. E. Green